


**A PHASE II STUDY OF DOSE-DENSE GEMCITABINE PLUS CISPLATIN (DDGC) IN PATIENTS WITH MUSCLE-INVASIVE BLADDER CANCER WITH BLADDER PRESERVATION FOR THOSE PATIENTS WHOSE TUMORS HARBOR DELETERIOUS DNA DAMAGE RESPONSE (DDR) GENE ALTERATIONS**

**Eligibility Criteria (see Section 3.0)**

- Histologically confirmed urothelial carcinoma of the bladder
- 10-20 unstained slides or 1 FFPE block from pre-treatment TUR available
- Clinical stage T2-T4aN0/xM0
- Candidate for radical cystectomy
- No prior systemic chemotherapy or radiation therapy for the bladder
- No major surgery or RT ≤ 4 weeks
- Non-pregnant and non-nursing
- Age ≥ 18 years
- ECOG PS = 0-1
- No comorbid conditions per Section 3.2.9

**Required Initial Laboratory Values**

- Absolute neutrophil count (ANC): ≥ 1000/mm<sup>3</sup>
- Platelet count: ≥ 100,000/mm<sup>3</sup>
- Calc. creatinine clearance: ≥ 55 mL/min
- Total bilirubin: ≤ 1.5 x ULN
- AST/ALT: ≤ 2.5 x ULN
- Alkaline phosphatase: ≤ 2.5 x ULN



Chemotherapy is to continue for 6 cycles or unacceptable adverse events (at least 4 cycles must be given for patients to proceed to Step 2 registration). Patients will be followed for five years after completion of chemotherapy or radical cystectomy, or until death, whichever comes first.

**Please refer to the full protocol text for a complete description of the eligibility criteria and treatment plan.**